

Fresubin® Renal

Available in Vanilla flavour
in 200 ml EasyBottle



Nutritional Information

Average content		100 ml	bottle = 200 ml
Energy value			
		kJ	1680
		(kcal)	(400)
Fat		g	17.8
of which saturated fatty acids		g	5.4
of which medium chain triglycerides		g	4.4
of which monounsaturated fatty acids		g	9.4
of which polyunsaturated fatty acids		g	2.94
of which EPA + DHA*		g	0.12
Carbohydrate		g	52.8
of which sugars		g	16.34
of which lactose		g	≤ 0.8
Fibre		g	2.4
Protein		g	6.0
Salt (Na x 2.5)		g	0.34
Water		ml	142
Osmolarity		mosmol/l	500
Minerals and trace elements			
Sodium	mg	68	136
Chloride	mg	62	64
Potassium	mg	100	200
Calcium	mg	84	168
Phosphorus	mg	55	110
Magnesium	mg	20	40
Iron	mg	2.0	4.0
Zinc	mg	1.8	3.6
Copper	µg	200	400
Iodine	µg	20	40
Selenium	µg	10	20
Manganese	mg	0.4	0.8
Chromium	µg	3.0	6.0
Molybdenum	µg	7.0	14.0
Fluoride	mg	0.3	0.6
Vitamins and other nutrients[^]			
Vitamin A	µg RE ^o	81.3	162.6
of which β-Carotene	µg RE ^o	33.3	66.6
Vitamin D ₃	µg	1.0	2.0
Vitamin E	mg α-TE ^{oo}	3.0	6.0
Vitamin K ₁	µg	10.5	21.0
Vitamin C	mg	10	20
Thiamin (vitamin B ₁)	mg	0.26	0.52
Riboflavin (vitamin B ₂)	mg	0.3	0.6
Vitamin B ₆	mg	0.67	1.34
Niacin	mg/mg NE ^{ooo}	3.4/4.2	6.8/8.4
Folic Acid	µg	100	200
Vitamin B ₁₂	µg	0.6	1.2
Pantothenic Acid	mg	1.2	2.4
Biotin	µg	12	24
Choline [^]	mg	55	110
Taurine [^]	mg	20	40
L-Carnitine [^]	mg	10	20
Inositol [^]	mg	24	48
Caloric distribution (energy%)			
Fat 40, carbohydrate 53, fibre 1.0, protein 6.0			
*eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA)			
^o retinol equivalents (RE), ^{oo} alpha-tocopherol equivalents (α-TE), ^{ooo} niacin equivalents (NE)			

General Information

Food for special medical purposes:

Nutritionally complete high-caloric (2 kcal/ml) oral nutritional supplement and tube feed, with fibre. Reduced protein (6 en%) and electrolytes. With fish oil. Carbohydrate modified with starch and isomaltulose. Clinically free from lactose, gluten free. For the dietary management of patients with or at risk of disease-related malnutrition, in particular with chronic kidney disease or non-catabolic forms of acute kidney injury.

Dosage:

To be determined by a health care professional according to patients' needs. Recommendation for supplementary nutrition 2-3 bottles (800-1200 kcal)/day and complete ≥ 5 bottles (2000 kcal)/day.

Important notes:

To be used under medical supervision. Monitor feeding rate when used for tube feeding. Suitable as sole source of nutrition. Not suitable for infants children < 3 years. Use with caution in children < 6 years. Not suitable for patients with galactosaemia. Ensure adequate fluid intake.

Not for parenteral (I.V.) use!

Instruction for use:

If used as tube feed: Recommended tube size is ≥ CH 8 for pump assisted feeding and for gravity feeding the minimum tube size is CH 8. Store at room temperature. Once opened, use within 24 hours. If used for sip feeding, shake well before use and drink slowly. If used for tube feeding, do not mix with drugs. Packaged in a protective atmosphere.

Ingredients

Water, maltodextrin, isomaltulose^o, vegetable oils (rapeseed oil, sunflower oil), tapioca dextrin, milk protein, medium chain triglycerides (MCT), emulsifiers (E 471, soya lecithins), flavouring, sodium citrate, fish oil (with soya lecithins), choline hydrogen tartrate, potassium chloride, potassium citrate, vit. C, magnesium oxide, myo-inositol, taurine, acidity regulator (E 524), l-carnitine, iron pyrophosphate, zinc sulphate, niacin, pantothenic acid, manganese chloride, vit. E, vit. B₆, sodium fluoride, copper sulphate, vit. B₂, vit. B₁, β-carotene, folic acid, vit. A, sodium selenite, biotin, potassium iodide, sodium molybdate, vit. K₁, chromium chloride, vit. B₁₂, vit. D₃.

^oisomaltulose is a source of glucose and fructose

Flavour

Vanilla

last updated: 09/2019



**FRESENIUS
KABI**

caring for life

Fresenius Kabi Deutschland GmbH
61346 Bad Homburg, Germany
Phone: +49 (0) 61 72 / 686-0
Enteral.nutrition@fresenius-kabi.com
www.fresenius-kabi.com