Diben® 1,5 kcal HP

Ready to use, in 500 ml or 1000 ml EasyBag



Nutritional Information

		100 m
Energy value	kJ	63
	(= kcal)	(150
Fat	g	7.0
of which saturated fatty acids	g	1.8
of which medium chain triglycerides	g	1.
of which monounsaturated fatty acids	g	3.6
of which polyunsaturated fatty acids	g	1.5
of which EPA* + DHA**	g	0.2
of which cholesterol	mg	≤ 1
Carbohydrate	g	13
of which starch	g	2.
of which sugars	g	7.
of which lactose	g	≤ 0.4
of which fructose	g	1.
of which isomaltulose	g	5.
Fibre	g	2.
Protein	g	7.
Salt (Na x 2.5)	g	0.1
Water	ml	7'
Osmolarity	mosmol/l	45
Minerals and trace elements		
Sodium	mg/mmol	55/2.
Chloride	mg/mmol	85/2.
Potassium	mg/mmol	180/4.
Calcium	mg/mmol	155/3.
Phosphorus	mg/mmol	110/3.
Magnesium	mg/mmol	30/1.
Iron	mg	2.
Zinc	mg	1.
Copper	μg	20
lodine	μg	2
Selenium	μg	1
Manganese		0.
	mg	
Chromium	μ g	2
Molybdenum	μg	1
Fluoride Vitamins and other nutrients^	mg	0.
Vitamin A	DE0	14
	μg RE°	
of which β-Carotene	μ g RE°	3.
Vitamin D ₃	μg	2.0
Vitamin E	mg α-TE°°	
Vitamin K ₁	μ g	1
Vitamin C	mg	1
Thiamin (vitamin B ₁)	mg	0.
Riboflavin (vitamin B ₂)	mg	0.2
Vitamin B ₆	mg	0.2
Niacin	mg/mg NE ^{oo}	° 2.4/4.
Folic Acid	μ g	4
Vitamin B ₁₂	μ g	0.
Pantothenic Acid	mg	0.
Biotin	μg	7.
Choline^	mg	5

Caloric distribution (energy%): Fat 42, carbohydrate 35, fibre 3, protein 20

General Information

Food for special medical purposes:Nutritionally complete high-caloric (1.5 kcal/ml), high-protein (20 energy%) tube feed with fibre. High in monounsaturated fatty acids, with MCT, with fish oil. Modified carbohydrate profile for improved glycaemic control. Clinically free from lactose, gluten free, low in sodium, low in cholesterol. For the dietary management of patients with or at risk of disease-related malnutrition in particular with impaired glucose metabolism such as diabetes mellitus and stress induced hyperglycaemia.

To be determined by a health care professional according to patients' needs. Recommendation for complete nutrition ≥ 1000 ml (1500 kcal)/day.

To be used under medical supervision. Monitor feeding rate. Suitable as sole source of nutrition. Not suitable for children < 3 years. Use with caution in children < 6 years. Not suitable for patients with galactosaemia. Ensure adequate fluid intake. Drug therapy for blood sugar control should be adjusted accordingly. Not for parenteral (I.V.) use!

Instructions for use:

Recommended tube size is ≥ CH 8 for pump assisted feeding and for gravity feeding the minimum tube size is CH 8. Store at room temperature. Once opened, use within 24 hours. Shake well before use! Do not use if bag is damaged or swollen or content is coagulated. Do not mix with drugs.

Ingredients

Water, <u>milk protein</u>, vegetable oils (sunflower oil, rapeseed oil, soya oil), isomaltulose^a, modified starch, maltodextrin, fructose, medium chain trigycerides (MCT), fish oil, inulin (from chicory), cellulose, potassium citrate, emulsifiers (E 471, <u>soya lecithin</u>), di-potassium hydrogen phosphate, sodium chloride, choline chloride, sodium citrate, stabilisers (E 460, E 466), magnesium oxide, vit. C, acidity regulator (E 330), iron pyrophosphate, zinc sulphate, niacin, manganese chloride, pantothenic acid, vit. E, copper sulphate, sodium fluoride, vit. B_2 , vit. B_6 , vit. B_1 , vit. A, β -carotene, chromium chloride, folic acid, sodium molybdate, sodium selenite, potassium iodide, biotin, vit. K₁, vit. D₃, vit. B₁₂

isomaltulose is a source of glucose and fructose

last updated: 09/2019



^{*}eicosapentaenoic acid (EPA), **docosahexaenic acid (DHA)
°retinol equivalents (RE), °° alpha-tocopherol equivalents (α-TE), °°° niacin equivalents (NE)